

# Chapter 39

### More About Matter Waves



#### **39.4: Two- and Three- Dimensional Electron Traps:**

The normalized wave function:



**Fig. 39-13** A rectangular corral—a two-dimensional version of the infinite potential well of Fig. 39-2—with widths 
$$L_x$$
 and  $L_y$ .

$$\psi_{nx, ny} = \sqrt{\frac{2}{L_x}} \sin\left(\frac{n_x \pi}{L} x\right) \sqrt{\frac{2}{L_y}} \sin\left(\frac{n_y \pi}{L} y\right),$$


The energy of the electron:

$$E_{nx,ny} = \left(\frac{h^2}{8mL_x^2}\right)n_x^2 + \left(\frac{h^2}{8mL_y^2}\right)n_y^2 = \frac{h^2}{8m}\left(\frac{n_x^2}{L_x^2} + \frac{n_y^2}{L_y^2}\right).$$

#### 39.4: Two- and Three- Dimensional Electron Traps, Rectangular Box

The energy of an electron trapped in a 3-D infinite potential box:

$$E_{nx,ny,nz} = \frac{h^2}{8m} \left( \frac{n_x^2}{L_x^2} + \frac{n_y^2}{L_y^2} + \frac{n_z^2}{L_z^2} \right).$$



**Fig. 39-14** A rectangular box—a three-dimensional version of the infinite potential well of Fig. 39-2—with widths  $L_x$ ,  $L_y$ , and  $L_z$ .



## Checkpoint 4

In the notation of Eq. 39-20, is  $E_{0,0}$ ,  $E_{1,0}$ ,  $E_{0,1}$ , or  $E_{1,1}$  the ground-state energy of an electron in a (two-dimensional) rectangular corral?

 $E_{1,1}$  (neither  $n_x$  nor  $n_y$  can be zero)

#### Example, Energy levels in a 2D infinite potential well:

An electron is trapped in a square corral that is a twodimensional infinite potential well (Fig. 39-13) with widths  $L_x = L_y$ .

(a) Find the energies of the lowest five possible energy levels for this trapped electron, and construct the corresponding energy-level diagram.

**Energy levels:** Because the well here is square, we can let the widths be  $L_x = L_y = L$ . Then Eq. 39-20 simplifies to

$$E_{nx,ny} = \frac{h^2}{8mL^2} (n_x^2 + n_y^2).$$
 (39-22)

The lowest energy states correspond to low values of the quantum numbers  $n_x$  and  $n_y$ , which are the positive integers 1, 2, ...,  $\infty$ . Substituting those integers for  $n_x$  and  $n_y$  in Eq. 39-22, starting with the lowest value 1, we can obtain the energy values as listed in Table 39-1. There we can see that

#### Table 39-1

#### **Energy Levels**

$n_x$	$n_y$	Energy <sup>a</sup>	$n_x$	$n_y$	Energy <sup>a</sup>
1	3	10	2	4	20
3	1	10	4	2	20
2	2	8	3	3	18
1	2	5	1	4	17
2	1	5	4	1	17
1	1	2	2	3	13
			3	2	13

<sup>&</sup>lt;sup>a</sup>In multiples of  $h^2/8mL^2$ .

These are the lowest five energy levels allowed the electron. Different quantum states may have the same energy.



13

 $E_{3,2}, E_{2,3}$ 

**Fig. 39-15** Energy-level diagram for an electron trapped in a square corral.

several of the pairs of quantum numbers  $(n_x, n_y)$  give the same energy. For example, the (1, 2) and (2, 1) states both have an energy of  $5(h^2/8mL^2)$ . Each such pair is associated with degenerate energy levels. Note also that, perhaps surprisingly,

From Table 39-1 (carefully keeping track of degenerate levels), we can construct the energy-level diagram of Fig. 39-15.

the (4,1) and (1,4) states have less energy than the (3,3) state.


#### Example, Energy levels in a 2D infinite potential well, cont.:

(b) As a multiple of  $h^2/8mL^2$ , what is the energy difference between the ground state and the third excited state?

**Energy difference:** From Fig. 39-15, we see that the ground state is the (1,1) state, with an energy of  $2(h^2/8mL^2)$ . We also see that the third excited state (the third state up from the ground state in the energy-level diagram) is the degenerate (1,3) and (3,1) states, with an energy of  $10(h^2/8mL^2)$ . Thus, the difference  $\Delta E$  between these two states is


$$\Delta E = 10 \left( \frac{h^2}{8mL^2} \right) - 2 \left( \frac{h^2}{8mL^2} \right) = 8 \left( \frac{h^2}{8mL^2} \right).$$
 (Answer)

These are the lowest five energy levels allowed the electron. Different quantum states may have the same energy.



**Fig. 39-15** Energy-level diagram for an electron trapped in a square corral.

#### 39.5: The Bohr Model of the Hydrogen Atom:



**Fig. 39-16** (a) Circular orbit of an electron in the Bohr model of the hydrogen atom. (b) The Coulomb force  $\vec{F}$  on the electron is directed radially inward toward the nucleus.

$$F = k \frac{|q_1||q_2|}{r^2},$$

$$-\frac{1}{4\pi\varepsilon_0}\frac{e^2}{r^2}=m\left(-\frac{v^2}{r}\right),$$

The angular momentum:  $\ell = rmv \sin \phi$ , =  $rmv \sin 90^{\circ}$ 

For quantization of *I*,  $rmv = n\hbar$ ,

$$v = \frac{n\hbar}{rm}$$
.

$$r = \frac{h^2 \varepsilon_0}{\pi m e^2} n^2, \quad \text{for } n = 1, 2, 3, \dots$$

$$r = an^{2}, \quad \text{for } n = 1, 2, 3, \dots,$$

$$a = \frac{h^{2} \varepsilon_{0}}{\pi m e^{2}} = 5.291772 \times 10^{-11} \text{ m} \approx 52.92 \text{ pm}.$$

#### 39.5: The Bohr Model of the Hydrogen Atom, Orbital energy is quantized:

$$E = K + U$$

$$= \frac{1}{2}mv^2 + \left(-\frac{1}{4\pi\varepsilon_0} \frac{e^2}{r}\right).$$

$$E = -\frac{1}{8\pi\epsilon_0} \frac{e^2}{r}.$$

$$E_n = -\frac{me^4}{8\varepsilon_0^2 h^2} \frac{1}{n^2}, \quad \text{for } n = 1, 2, 3, \dots,$$

$$E_n = -\frac{2.180 \times 10^{-18} \,\text{J}}{n^2} = -\frac{13.61 \,\text{eV}}{n^2}, \quad \text{for } n = 1, 2, 3, \dots$$

#### 39.5: The Bohr Model of the Hydrogen Atom, Energy Changes:

$$hf = \Delta E = E_{\text{high}} - E_{\text{low}}.$$



$$\frac{1}{\lambda} = -\frac{me^4}{8\varepsilon_0^2 h^3 c} \left( \frac{1}{n_{\text{high}}^2} - \frac{1}{n_{\text{low}}^2} \right).$$



$$\frac{1}{\lambda} = R \left( \frac{1}{n_{\text{low}}^2} - \frac{1}{n_{\text{high}}^2} \right),$$

$$R = \frac{me^4}{8\varepsilon_0^2 h^3 c} = 1.097\ 373 \times 10^7\ \text{m}^{-1}$$
 (Rydberg Constant)

#### 39.9: Schrodinger's Equation and The Hydrogen Atom:





Fig. 39-17 The potential energy U of a hydrogen atom as a function of the separation r between the electron and the central proton. The plot is shown twice (on the left and on the right) to suggest the three-dimensional spherically symmetric trap in which the electron is confined.


#### Schrodinger's Equation and The Hydrogen Atom:



**Fig. 39-18** (a) An energy-level diagram for the hydrogen atom. Some of the transitions for (b) the Lyman series. For each, the longest four wavelengths and the series-limit wavelength are plotted on a wavelength axis. Any wavelength shorter than the series-limit wavelength is allowed.

#### Schrodinger's Equation and The Hydrogen Atom:

**Fig. 39-18** Some of the transitions for (c) the Balmer series, and (d) the Paschen series. For each, the longest four wavelengths and the series-limit wavelength are plotted on a wavelength axis. Any wavelength shorter than the series-limit wavelength is allowed.



#### **Quantum Numbers and The Hydrogen Atom:**

Each set of quantum numbers  $(n, l, m_l)$  identifies the wave function of a particular quantum state. The quantum number n, is called the **principal quantum number.** The **orbital quantum number** l is a measure of the magnitude of the angular momentum associated with the quantum state. The **orbital magnetic quantum number**  $m_l$  is related to the orientation in space of this angular momentum vector.

The restrictions on the values of the quantum numbers for the hydrogen atom, as listed in Table 39-2, are not arbitrary but come out of the solution to Schrödinger's equation.

Table 39-2					
Quantum Numbers for the Hydrogen Atom					
Symbol	Name	Allowed Values			
n	Principal quantum number	1, 2, 3,			
$\ell$	Orbital quantum number	$0,1,2,\ldots,n-1$			
$m_\ell$	Orbital magnetic quantum number	$-\ell, -(\ell-1), \ldots, +(\ell-1), +\ell$			